



Anxiety and Fear of Recurrence in Cancer Survivors

Dr. Ioanna Tsatsou*

Cancer Nurse, Msc, PhD, Oncology-Hematology Department, Hellenic Airforce General Hospital, Athens, Greece



WebLog Open Access Publications
Article ID : wjda.2025.e0504
Author : Dr. Ioanna Tsatsou, Ph.D.

Abstract

Cancer survivorship has significantly increased due to advancements in early detection and treatment. However, survivors often face psychological challenges, notably anxiety and fear of cancer recurrence (FCR). These concerns can persist long after treatment, adversely affecting quality of life (QoL). This narrative review explores the prevalence, implications, assessment methods, and management strategies for anxiety and FCR among cancer survivors. Recent studies indicate that over half of survivors experience moderate to high levels of FCR, with younger age and female gender being significant risk factors. Persistent anxiety and FCR are associated with increased depression, reduced QoL, and impaired daily functioning. Effective management requires a multidisciplinary approach, integrating psychological interventions and supportive care. Addressing these psychological concerns is crucial for improving survivorship outcomes.

Keywords: Cancer Survivorship, Anxiety, Fear of Cancer Recurrence, Quality of Life, Psychological Assessment, FCRI

Introduction

The population of cancer survivors has been consistently rising, attributed to advancements in cancer detection and treatment methods. While this is a positive development, it brings to light the long-term psychological challenges faced by survivors. Among these, anxiety and fear of cancer recurrence (FCR) are predominant concerns that can significantly impact a survivor's quality of life (QoL). Understanding the prevalence, implications, and management of these psychological issues is essential for comprehensive survivorship care [1].

Anxiety in Cancer Survivors

Anxiety is a common psychological response among cancer survivors, often stemming from uncertainties about the future, fear of recurrence, and the aftermath of treatment. Studies have shown that anxiety can persist long after treatment completion, affecting daily functioning and overall well-being. For instance, nearly 42% of breast cancer survivors experience anxiety, which can be exacerbated by factors such as undergoing mastectomy or being diagnosed at a younger age [2]. The manifestations of anxiety in cancer survivors can range from mild worry to severe panic attacks, often interfering with sleep, concentration, and social interactions. Persistent anxiety can also hinder adherence to follow-up care and surveillance, potentially impacting long-term health outcomes [3].

Fear of Cancer Recurrence (FCR)

FCR refers to the fear, worry, or concern that cancer could return or progress. It is a prevalent and distressing issue among survivors, often leading to heightened vigilance, frequent medical consultations, and avoidance behaviors. FCR can be distinguished from general anxiety by its specific focus on cancer returning. It is influenced by factors such as age, gender, type of cancer, and time since diagnosis. Younger survivors and female individuals tend to report elevated levels of FCR [1, 4]. The fear of recurrence can be so overwhelming that it affects a survivor's ability to enjoy life, plan for the future, and engage in meaningful activities. This persistent fear can lead to chronic stress, depression, and a diminished sense of well-being [5].

Prevalence of Anxiety and FCR in Cancer Survivors

Recent studies highlight the significant prevalence of anxiety and FCR among cancer survivors. A systematic review and meta-analysis involving over 9,000 participants found that approximately 59% experienced at least moderate levels of FCR, with about 19% reporting high levels [1]. Similarly, anxiety remains a persistent issue, with factors such as younger age, female gender, and certain cancer types (e.g., lung and melanoma) associated with higher anxiety and FCR levels [3]. These

OPEN ACCESS

*Correspondence:

Dr. Ioanna Tsatsou, Cancer Nurse,
Msc, PhD, Oncology-Hematology
Department, Hellenic Airforce General
Hospital, Athens, Greece,
E-mail: itsatsou@uniwa.gr

Received Date: 29 Apr 2025

Accepted Date: 02 May 2025

Published Date: 05 May 2025

Citation:

Ioanna Tsatsou. Anxiety and Fear
of Recurrence in Cancer Survivors.
WebLog J Depress Anxiety. wjda.2025.
e0504. [https://doi.org/10.5281/
zenodo.15981188](https://doi.org/10.5281/zenodo.15981188)

Copyright© 2025 Dr. Ioanna

Tsatsou. This is an open access
article distributed under the Creative
Commons Attribution License, which
permits unrestricted use, distribution,
and reproduction in any medium,
provided the original work is properly
cited.

statistics underscore the need for routine psychological assessment and support as integral components of survivorship care [4].

Implications of Anxiety and FCR on Quality of Life

Anxiety and FCR significantly impact the QoL of cancer survivors. They are associated with increased risk of depression, reduced emotional and social functioning, and impaired daily activities. In long-term breast cancer survivors, higher FCR levels correlate with increased depression and decreased QoL [6]. Persistent psychological distress can also lead to increased healthcare utilization and hindered return to normal life roles [7]. Moreover, the chronic stress associated with anxiety and FCR can have physiological effects, potentially impacting immune function and overall health. Addressing these psychological concerns is therefore not only essential for mental well-being but also for physical health and recovery [5].

Assessment of Anxiety and FCR

Effective assessment of anxiety and FCR is crucial for identifying survivors in need of support. The Fear of Cancer Recurrence Inventory (FCRI) is a validated tool commonly used to measure FCR levels. The short form (FCRI-SF) consists of 9 items, with scores indicating the severity of FCR. A score of 22 or higher suggests clinical levels of FCR requiring intervention [8]. Additionally, structured interviews and questionnaires like the General Anxiety Disorder-7 (GAD-7) can assess general anxiety levels [4]. Regular screening using these tools can facilitate early identification and timely intervention, potentially mitigating the long-term impact of anxiety and FCR on survivors' lives [9].

Management Strategies

Addressing anxiety and FCR in cancer survivors requires a multidisciplinary approach. Psychological interventions, particularly cognitive-behavioral therapy (CBT), have been shown to be effective in reducing symptoms. CBT assists individuals in recognizing and confronting unhelpful thoughts and behaviors, promoting the development of more effective coping strategies [5]. Additionally, mindfulness-based stress reduction (MBSR) programs can aid in managing anxiety and improving QoL [3, 10].

Pharmacological treatments, such as antidepressants or anxiolytics, may be considered for individuals with severe symptoms. However, these should be prescribed cautiously, considering potential side effects and interactions with ongoing cancer treatments [6].

Support groups and peer counseling can also provide emotional support, reduce feelings of isolation, and offer practical coping strategies. Engaging in regular physical activity, maintaining a healthy lifestyle, and fostering strong social connections are additional measures that can alleviate anxiety and FCR [7].

Conclusion

Anxiety and fear of cancer recurrence are prevalent and impactful concerns among cancer survivors. They significantly affect emotional well-being, daily functioning, and overall quality of life. Regular

assessment using validated tools is essential for early identification. Integrating psychological support into survivorship care plans can help address these issues, improving outcomes and QoL for cancer survivors.

References

- Luigjes-Huizer YL, Tauber NM, Humphris G, Kasparian NA, Lam WWT, Lebel S, Simard S, Smith AB, Zachariae R, Afyanti Y, Bell KJL, Custers JAE, de Wit NJ, Fisher PL, Galica J, Garland SN, Helsper CW, Jeppesen MM, Liu J, Mititelu R, Monninkhof EM, Russell L, Savard J, Speckens AEM, van Helmondt SJ, Vatandoust S, Zdenkowski N, van der Lee ML. What is the prevalence of fear of cancer recurrence in cancer survivors and patients? A systematic review and individual participant data meta-analysis. *Psychooncology*. 2022 Jun; 31(6): 879-892.
- Gnall KE, Emrich M, Magin ZE, Park CL, Bellizzi KM, Sanft T. Anxiety and fear of cancer recurrence as predictors of subsequent pain interference in early cancer survivorship: Exploring the moderating roles of cognitive and emotional factors. *J Behav Med*. 2024 Dec; 47(6): 980-993.
- Simard S, Thewes B, Humphris G, Dixon M, Hayden C, Mireskandari S, Ozakinci G. Fear of cancer recurrence in adult cancer survivors: a systematic review of quantitative studies. *J Cancer Surviv*. 2013 Sep; 7(3): 300-22.
- Simard S, Savard J. Fear of Cancer Recurrence Inventory: development and initial validation of a multidimensional measure of fear of cancer recurrence. *Support Care Cancer*. 2009; 17(3): 241-251.
- Smith AB, Costa D, Galica J, Lebel S, Tauber N, van Helmondt SJ, Zachariae R. Spotlight on the Fear of Cancer Recurrence Inventory (FCRI). *Psychol Res Behav Manag*. 2020 Dec 21; 13: 1257-1268.
- Tran TXM, Jung SY, Lee EG, Cho H, Kim NY, Shim S, Kim HY, Kang D, Cho J, Lee E, Chang YJ, Cho H. Fear of Cancer Recurrence and Its Negative Impact on Health-Related Quality of Life in Long-term Breast Cancer Survivors. *Cancer Res Treat*. 2022 Oct; 54(4): 1065-1073.
- Lebel S, Simard S, Harris C, Feldstain A, Beattie S, McCallum M, Lefebvre M, Savard J, Devins GM. Empirical validation of the English version of the Fear of Cancer Recurrence Inventory. *Qual Life Res*. 2016 Feb; 25(2): 311-321.
- Costa DS, Dieng M, Cust AE, Butow PN, Kasparian NA. Psychometric properties of the Fear of Cancer Recurrence Inventory: an item response theory approach. *Psychooncology*. 2016 Jul; 25(7): 832-838.
- Thewes B, Bell ML, Butow P. Fear of cancer recurrence in young early-stage breast cancer survivors: the role of metacognitive style and disease-related factors. *Psychooncology*. 2013 Sep; 22(9): 2059-2063.
- Reich RR, Lengacher CA, Alinat CB, Kip KE, Paterson C, Ramesar S, Han HS, Ismail-Khan R, Johnson-Mallard V, Moscoso M, Budhrani-Shani P, Shivers S, Cox CE, Goodman M, Park J. Mindfulness-Based Stress Reduction in Post-treatment Breast Cancer Patients: Immediate and Sustained Effects Across Multiple Symptom Clusters. *J Pain Symptom Manage*. 2017 Jan; 53(1): 85-95.