



Treatment and Interventions for Attention-Deficit Hyperactivity Disorder: Best Practices and New Directions

Dr. Shungui Xiang¹, Dr. Yin Tang², Dr. Lei Zhao³ and Dr. Jun Zhang^{4,5*}

¹Library, Tourism College of Zhejiang, Hangzhou, 311231, China

²School of Travel Services and Management, Tourism College of Zhejiang, Hangzhou, 311231, China

³Faculty of Applied Law, Zhejiang Police Vocational Academy, Hangzhou, 310018, China

⁴School of Business Administration, Tourism College of Zhejiang, Hangzhou, 311231, China

⁵College of Education, Sehan University, Jeollanam-do, 650106, Republic of Korea



WebLog Open Access Publications

Article ID : wjpbs.2025.g2901
Author : Dr. Jun Zhang, Ph.D.

Abstract

Attention-Deficit/Hyperactivity Disorder (ADHD) is a common neurodevelopmental disorder that affects children and can persist into adulthood, posing significant challenges in academic, social, and family settings. Pharmacological treatment remains one of the most widely used and effective approaches. In addition to medication, behavioral therapy has been proven effective in improving behavior and emotional regulation, and is often combined with medication for a more comprehensive treatment. Finally, we highlight future directions in ADHD treatment, focusing on more personalized and less invasive approaches, as well as the potential role of digital health tools in managing ADHD.

Keywords: Attention-Deficit Hyperactivity Disorder; Medication Treatment; Behavioral Therapy; Personalized Treatment

Introduction

Attention-Deficit/Hyperactivity Disorder (ADHD) is a common childhood neurodevelopmental disorder with a relatively high prevalence rate. It is characterized by inattention, hyperactivity, and impulsive behaviors, which significantly affect children's learning, social interactions, and family life. ADHD can impair academic performance, as affected children often struggle to concentrate in class and have difficulty completing assignments given by teachers. In more severe cases, it may lead to poor peer relationships, underdeveloped social skills, and even social isolation or rejection. Behaviorally, these children often exhibit frequent impulsivity and emotional instability, which can trigger family conflicts and negatively impact the mental well-being of family members [2]. Without timely and effective intervention, symptoms may persist into adolescence or adulthood, potentially resulting in emotional disorders, depression, anxiety, and even affecting career development and family life [7].

Symptoms and Types of ADHD

According to the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), symptoms of Attention-Deficit/Hyperactivity Disorder (ADHD) must appear before the age of 12 and persist for more than six months in multiple settings, such as at home and school [1]. The symptoms of ADHD are categorized into three types: predominantly inattentive type, predominantly hyperactive-impulsive type, and combined type [1]. Children with the inattentive type often struggle to maintain focus for extended periods, are easily distracted, and have difficulty completing tasks or organizing activities in a structured manner [2]. They may overlook details, work carelessly, frequently lose items, or forget important tasks. Additionally, these children may show a lack of interest in daily activities, leading to inefficiency in both academic and family environments. Children with the hyperactive-impulsive type find it hard to remain seated or stay calm. They often move constantly, may climb on furniture or run around excessively—even in inappropriate settings [8]. The combined type includes symptoms of both inattention and hyperactivity-impulsivity, and is the most common form of ADHD. Children with this type often face significant challenges in multiple areas, including academic performance, behavior regulation, and social adaptation [2].

It is worth noting that ADHD symptoms can vary significantly between individuals, and some

OPEN ACCESS

*Correspondence:

Dr. Jun Zhang, Department of Education, Sehan University, Yeongam-gun, Jeollanam-do, 650106, Republic of Korea,

E-mail: zhangjunahnu@163.com

Received Date: 16 Jun 2025

Accepted Date: 28 Jul 2025

Published Date: 29 Jul 2025

Citation:

Xiang S, Tang Y, Zhao L, Zhang J. Treatment and Interventions for Attention-Deficit Hyperactivity Disorder: Best Practices and New Directions. *WebLog J Psychiatry Behav Sci.* wjpbs.2025.g2901. <https://doi.org/10.5281/zenodo.16558885>

Copyright© 2025 Dr. Jun Zhang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

symptoms may improve with age. However, even if symptoms diminish over time, ADHD may continue to affect adolescents and adults, particularly in areas such as work, family life, and social relationships [5].

Etiology and Risk Factors of ADHD

Attention-Deficit/Hyperactivity Disorder (ADHD) is a neurodevelopmental disorder, the exact etiology of which remains not fully understood. However, research suggests that the onset of ADHD is the result of the interaction of multiple genetic, environmental, and neurobiological factors. Among them, genetic factors are considered one of the most significant causes of ADHD. Variations in the dopamine transporter gene (DAT1) and the dopamine receptor gene (DRD4) have been associated with the development of ADHD [2]. These genetic mutations may affect neurotransmission and executive function in the brain, leading to the manifestation of ADHD symptoms.

Maternal health during pregnancy may also increase the risk of ADHD. Factors such as maternal smoking, alcohol consumption, or exposure to toxic substances during pregnancy can adversely affect fetal brain development and thus raise the likelihood of ADHD [1]. These prenatal influences may result in abnormalities in brain structure and function, which can impair attention regulation and behavioral control [10].

Modern neuroimaging studies have shown that children with ADHD often exhibit reduced activity or structural abnormalities in brain regions such as the prefrontal cortex, basal ganglia, and cerebellum. This suggests that these regions play a crucial role in the development of ADHD [2]. These neurobiological abnormalities are thought to be associated with dysregulation of neurotransmitters like dopamine, affecting the brain's self-regulation mechanisms and leading to deficits in attention and behavior.

Treatment and Intervention for ADHD

In clinical practice, there are various treatment and intervention approaches for Attention-Deficit/Hyperactivity Disorder (ADHD), including medication, behavioral therapy, and combined therapy. These treatments aim to alleviate symptoms, improve functional impairments, and help individuals adapt to academic, social, and family life.

Medication is currently one of the most common treatment methods for ADHD. Pharmacological treatments primarily include stimulant medications (such as methylphenidate and amphetamines) and non-stimulant medications (such as atomoxetine and guanfacine). Stimulant medications work by enhancing the activity of dopamine and norepinephrine in the brain to improve attention and impulse control, while non-stimulant medications regulate neurotransmitter levels through different mechanisms and tend to have fewer side effects [1]. These medications can significantly improve attention, behavioral control, and academic performance in children with ADHD. However, medication treatment may be accompanied by side effects such as decreased appetite and sleep disturbances. Behavioral therapy aims to help individuals with ADHD improve their adaptive functioning in daily life by reinforcing positive behaviors and reducing maladaptive ones. Common behavioral techniques include reward systems, time management training, and organizational skills training [2]. By establishing clear rules, reward mechanisms, and structured behavioral management strategies, behavioral therapy can

effectively improve the behavior of children with ADHD at school and at home. A combination of medication and behavioral therapy is often considered the most effective treatment approach. Research has shown that the integration of pharmacological and behavioral interventions can more comprehensively improve the symptoms and functional outcomes of children with ADHD [6]. Furthermore, as children with ADHD grow older, treatment strategies in adulthood may need to be adjusted to focus more on emotional regulation, vocational adaptation, and the development of social skills. Overall, the treatment and intervention of ADHD should be individualized based on the specific symptoms, age, living environment, and level of family support for each patient.

Conclusion

Attention-Deficit/Hyperactivity Disorder (ADHD) poses significant challenges for both patients and their families. While medication remains one of the primary treatment methods for ADHD, it also presents certain challenges and limitations. For example, medications may cause side effects such as reduced appetite, sleep disturbances, and mood fluctuations. This highlights the need for future research to focus on developing more precise and individualized pharmacological treatments, aiming to discover medications with fewer side effects and longer-lasting therapeutic effects.

References

1. American Psychiatric Association, DSM-5 Task Force. Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.). American Psychiatric Publishing, Inc. 2013.
2. Barkley R.A. (Ed.). Attention-deficit hyperactivity disorder: A handbook for diagnosis and treatment (4th ed.). The Guilford Press. 2015.
3. Huang Y.X. A brief analysis of the education and management of students with ADHD. *Time Education*, 2021; (8), 23-24.
4. Liang W. Education for attention-deficit hyperactivity disorder. *Journal of Taiyuan Urban Vocational Technical College*, 2013; (6), 128-128.
5. MTA Cooperative Group. A 14-month randomized clinical trial of treatment strategies for Attention-Deficit Hyperactivity Disorder. *Archives of General Psychiatry*, 1999; 56(12), 1073-1086. <https://doi.org/10.1001/archpsyc.56.12.1073>
6. Pelham W.E, Burrows-MacLean L, Gnagy E.M, Fabiano G.A, Coles E.K, Wymbs B.T, et al. A dose-ranging study of behavioral and pharmacological treatment in social settings for children with ADHD. *Journal of Abnormal Child Psychology*, 2014; 42(6), 1019-1031. <https://doi.org/10.1007/s10802-013-9843-8>
7. Perroud N, Cordera P, Zimmermann J, Michalopoulos G, Bancila V, Prada P, et al. Comorbidity between attention deficit hyperactivity disorder (ADHD) and bipolar disorder in a specialized mood disorders outpatient clinic. *Journal of Affective Disorders*, 2014; 168, 161-166. doi: 10.1016/j.jad.2014.06.053.
8. Tucha O, Walitza S, Mecklinger L, et al. Attentional functioning in children with ADHD – predominantly hyperactive-impulsive type and children with ADHD – combined type. *Journal of Neural Transmission*, 2006; 113, 1943-1953. <https://doi.org/10.1007/s00702-006-0496-4>
9. Wang Y. Emphasizing medication treatment for ADHD. *Jiangsu Health Care*, 2022; (12), 10-11.
10. Zeeuw P.D, Zwart F, Schrama R, Engeland H.V & Durston S. Prenatal exposure to cigarette smoke or alcohol and cerebellum volume in Attention-Deficit Hyperactivity Disorder and typical development. *Translational Psychiatry*, 2012; 2, e84. <https://doi.org/10.1038/tp.2012.12>