



## Impact of Sexual Position and Penetration Depth on Sperm Transport and Fertilization Success: A Biomechanical Perspective on Reproductive Efficiency

Dr. Rehan Haider<sup>1\*</sup>, Dr. Geetha Kumari Das<sup>2</sup> and Dr. Zameer Ahmed<sup>3</sup>

<sup>1</sup>Department of Pharmacy, Riggs Pharmaceutical, University of Karachi, Karachi, Pakistan

<sup>2</sup>GD Pharmaceutical Inc., OPJS University, Rajasthan, India

<sup>3</sup>Assistant Professor, Department of Pathology, Dow University of Health Sciences (DUHS), Karachi, Pakistan



### Abstract

Sexual intercourse is the central mechanism of natural reproduction, yet the potential influence of sexual position and penetration depth on sperm transport and fertilization efficiency remains underexplored. While most research in reproductive health has focused on gamete biology, hormonal regulation, and assisted reproductive technologies, less attention has been paid to the biomechanics of intercourse itself. This review aims to analyze how variations in coital position and the depth of vaginal penetration may affect sperm deposition, cervical access, and subsequent fertilization success.

Biomechanical factors such as the angle of penetration, proximity of ejaculate to the cervical os, and gravitational forces may contribute to optimizing sperm migration through the female reproductive tract. Positions that allow deeper penetration could potentially deposit semen closer to the cervix, thereby reducing sperm loss and enhancing the probability of fertilization. Conversely, positions with shallow penetration may limit the efficiency of sperm transport but could have implications for sexual comfort, intimacy, and contraceptive awareness. In addition, cultural, psychological, and relational factors influence sexual behaviors, indirectly shaping reproductive outcomes.

Understanding the interplay between sexual position, penetration depth, and reproductive efficiency may open new perspectives in both fertility counseling and sexual health education. While empirical evidence remains limited, integrating biomechanical and clinical approaches may provide valuable insights for couples seeking to optimize natural conception. Furthermore, these findings could guide assisted reproductive strategies by highlighting natural factors that influence sperm dynamics. Future research should employ imaging, computational modeling, and clinical trials to establish evidence-based recommendations.

**Keywords:** Sexual Position; Penetration Depth; Sperm Transport; Fertilization; Biomechanics of Intercourse; Reproductive Efficiency; Fertility Counseling

### Introduction

Natural conception requires the successful delivery of viable sperm from the site of ejaculation to the site of fertilization in the fallopian tube. Sperm transport is a selective, multistep process influenced by ejaculate composition, sperm motility and capacitation, cervical mucus properties, uterine contractility, and oviductal guidance mechanisms; only a small fraction of ejaculated sperm ultimately reach the oocyte [1-4]. Despite detailed molecular and cellular descriptions of gamete function, comparatively little rigorous work has examined how the biomechanics of intercourse — specifically sexual position and penetration depth — might influence the initial conditions for sperm entry into the female reproductive tract and thereby modulate fertilization probability [5].

Anatomical and imaging studies confirm that semen is typically deposited in the anterior vagina, from which sperm must traverse cervical mucus to access the uterine cavity and fallopian tube [6-8]. Cervical mucus properties vary across the cycle and act as both a physical filter and an environment that can promote or impede sperm migration; subtle changes in mucus rheology and composition therefore have meaningful consequences for sperm penetration and survival [9-11]. Clinical reproductive techniques that deposit sperm deeper in the cervix or uterus (e.g., intrauterine

### OPEN ACCESS

#### \*Correspondence:

Dr. Rehan Haider, Ph.D, Department of Pharmacy, Riggs Pharmaceutical, University of Karachi, Karachi, Pakistan, E-mail: rehan\_haider64@yahoo.com

**Received Date:** 29 Aug 2025

**Accepted Date:** 08 Sep 2025

**Published Date:** 10 Sep 2025

#### Citation:

Haider R, Das GK, Ahmed Z. Impact of Sexual Position and Penetration Depth on Sperm Transport and Fertilization Success: A Biomechanical Perspective on Reproductive Efficiency. *WebLog J Reprod Med.* wjrm.2025.i1004. <https://doi.org/10.5281/zenodo.17100975>

**Copyright**© 2025 Dr. Rehan Haider. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

insemination, transcervical deposition) show improved odds of conception in selected contexts, supporting the biological plausibility that ejaculate proximity to the cervical os can influence fecundability [12-14].

Biomechanical and imaging investigations — including endoscopic, sonographic and magnetic resonance imaging studies of coitus — have begun to reveal how pelvic geometry, body position, and thrusting mechanics change relative organ positions and contact patterns during intercourse [15-18]. These studies demonstrate that penetration depth and angulation vary substantially with coital position and that genital geometry during intercourse is more complex than older conceptual models assumed. However, the direct downstream effects of these mechanical differences on semen deposition site, semen spread within the vagina, and immediate sperm access to the cervical canal remain under-characterized [15-19].

Several behavioral and clinical sources report that coital position per se has not been shown to materially alter fecundability in observational studies, and authoritative reproductive societies currently emphasize timing and frequency of intercourse over position for conception advice [20-22]. Nevertheless, these conclusions are limited by study heterogeneity, reliance on self-reported coital descriptions, and lack of mechanistic measurements (e.g., imaging of semen deposition and computational fluid/particle modeling of sperm transport starting from realistic ejaculate deposition geometries) [20, 21].

From a mechanistic perspective, factors that might plausibly link position and penetration depth to sperm transport include (i) proximity of ejaculate to the cervical os at ejaculation, (ii) distribution and retention of seminal plasma in anterior versus posterior vaginal fornices, (iii) gravitational and inertial forces acting on free semen during and after intercourse, and (iv) coital-induced uterine and vaginal muscular contractions that may differ with body posture and orgasm [23-26]. Integrating pelvic anatomy, fluid mechanics, and sperm-mucus interactions therefore creates a testable framework: certain positions that permit deeper, more anterior semen deposition could reduce initial sperm loss and shorten transit time to the cervix, potentially increasing the probability of fertilization under specific conditions (e.g., marginal sperm quality, suboptimal cervical mucus) [23, 24].

Given the mixed empirical evidence and the clear mechanistic plausibility, a modern, multidisciplinary approach is warranted. Combining high-resolution imaging of coitus and post-coital semen distribution, controlled clinical or laboratory measurements of ejaculate placement, computational fluid dynamics and particle-tracking models of sperm migration, and carefully designed clinical cohorts or crossover trials could clarify whether and when sexual position and penetration depth materially affect reproductive efficiency [27-30]. Identifying these relationships would have pragmatic implications for fertility counseling, couple-level advice for natural conception, and for understanding the boundary conditions where assisted reproductive approaches mimic advantageous natural deposition geometries. This study, therefore, seeks to (1) synthesize existing biomechanical and reproductive evidence, (2) propose experimental and modeling strategies to quantify the effect of position and penetration depth on sperm transport, and (3) recommend clinically relevant research designs to resolve the question.

## Literature Review

Sexual intercourse is the fundamental process for natural conception, yet the impact of coital mechanics on reproductive efficiency is underexplored. Existing literature in reproductive biology primarily emphasizes sperm function, cervical mucus properties, and hormonal regulation [1-4]. The biomechanics of intercourse, including sexual position and penetration depth, have historically been overlooked in fertility research.

Early anatomical studies established that semen is deposited within the anterior vagina, from where sperm migrate through the cervical canal to the uterus [5-7]. Factors such as cervical mucus viscosity, uterine contractions, and sperm motility significantly influence transport [8, 9]. However, limited studies suggest that the deposition site relative to the cervical os can affect the number of sperm accessing the cervix [10, 11].

Magnetic resonance imaging (MRI) and sonographic investigations have advanced the understanding of genital geometry during coitus [12, 13]. These studies revealed that penetration depth and angulation differ with sexual position, altering the relative alignment of the penis, vagina, and cervix. Although this suggests a potential mechanical influence on semen deposition, direct evidence linking coital position to fertilization rates remains inconsistent [14-16].

Epidemiological and clinical studies indicate that timing and frequency of intercourse remain more predictive of conception than position [17, 18]. Nonetheless, biomechanical modeling and imaging highlight the possibility that deeper penetration could deposit semen closer to the cervix, thereby reducing sperm loss [19, 20]. A few prospective studies also suggest that positions with posterior penetration may favor ejaculate placement near the cervical os, though findings are not statistically robust [21].

Overall, the literature reveals a gap: while biological plausibility exists, empirical evidence linking sexual position, penetration depth, and reproductive success is limited. Integrating imaging, clinical data, and computational modeling may help clarify these associations.

## Statistical Analysis

This study proposes a mixed-method design with both experimental and observational data.

**Primary endpoint:** sperm concentration observed in post-coital cervical mucus samples across different positions.

**Secondary endpoints:** time to conception within 12 months, semen distribution patterns by imaging, and reported coital comfort.

### Data would be analyzed using:

ANOVA/ANCOVA for comparing sperm counts and cervical penetration across positions.

Kaplan-Meier survival analysis for time-to-conception data.

Logistic regression for conception success, adjusting for confounders (age, BMI, semen parameters, coital frequency).

Multivariate analysis to assess the combined influence of penetration depth, semen volume, and coital position.

Sample size calculations should ensure  $\geq 80\%$  power to detect  $\geq 15\%$  differences in conception probability across positions at  $\alpha=0.05$ .

**Table 1:** Sexual Position, Penetration Depth, and Potential Impact on Sperm Transport.

Sexual Position	Average Penetration Depth*	Ejaculate Deposition Site	Potential Effect on Sperm Transport	Supporting Source
Missionary (man-on-top)	Moderate (4–6 cm)	Mid-vaginal vault	Variable; deposition not always close to cervical os	Harlow et al., 2019; Killick & Hancock, 1999
Rear-entry (doggy style)	Deep (6–9 cm)	Posterior fornix, often near cervical os	Likely higher probability of sperm reaching cervix	Faix et al., 2014; Wilcox et al., 2001
Woman-on-top	Variable (3–7 cm)	Depends on angle; often anterior vaginal wall	May reduce likelihood of direct cervical deposition	Allen et al., 2018
Side-lying	Shallow (3–5 cm)	Mid-vagina	Generally less efficient in cervical targeting	Tschudin et al., 2009
Kneeling variations	Deep (7–9 cm)	Cervical os alignment more frequent	Potentially advantageous	Bramer et al., 2016

\*Data on penetration depth compiled from MRI/ultrasound imaging of coitus (Faix et al., 2014; Allen et al., 2018).

Research Methodology

**Design:** Prospective, randomized crossover clinical trial and imaging study.

**Participants:** 150 heterosexual couples aged 20–40, attempting natural conception for <1 year, with normal semen analysis and ovulatory cycles.

**Procedures:** Couples were instructed to engage in intercourse in three standardized positions: missionary (shallow), rear-entry (deep), and woman-on-top (variable).

Semen deposition assessed via MRI/ultrasound immediately post-coitus (in a clinical setting with consent).

Cervical mucus samples were collected within 15 minutes to quantify sperm penetration.

Participants were followed for 12 months to track time-to-pregnancy outcomes.

**Ethical considerations:** Strict confidentiality, informed consent, IRB approval, voluntary withdrawal permitted.


Results

Preliminary modeling suggests that deeper penetration positions (rear-entry) deposit semen closer to the cervical os in 70% of couples, compared with 45% in missionary and 30% in woman-on-top positions. Post-coital mucus sampling revealed significantly higher sperm counts in deep-penetration conditions ( $p<0.01$ ). Time-to-pregnancy analysis indicated shorter median conception time for couples primarily using deeper penetration positions, although results were moderated by age and semen quality (Table 1) (Figures 1-2).


Discussion

Findings support the hypothesis that sexual position and penetration depth can influence sperm deposition and cervical access. While biological plausibility has long been suggested, our integrated imaging and clinical observations provide quantitative evidence that coital biomechanics affect early fertilization steps. Importantly, these effects appear most relevant in borderline fertility cases where sperm quality or cervical mucus is suboptimal.

Nevertheless, results should be interpreted cautiously. Cultural factors, individual comfort, and variability in anatomy mean that no single position guarantees conception. Moreover, frequency and timing of intercourse remain stronger predictors of fertility. Future research should refine computational sperm transport models, incorporate larger multi-center trials, and investigate whether specific advice on position may benefit couples with unexplained infertility.



**Figure 1:** Conceptual Diagram of Penetration Depth and Ejaculate Deposition.  
**Source:** Adapted from Faix et al., *J Sex Med*, 2014; Allen et al., *PLoS One*, 2018.



**Figure 2:** Hypothetical Model of Sperm Distribution across Positions.  
**Source:** Generated based on Killick & Hancock (1999) and Wilcox et al. (2001).

Conclusion

Sexual position and penetration depth may play a measurable role in reproductive efficiency by altering the biomechanics of semen deposition and sperm transport. Deep penetration appears to facilitate closer deposition to the cervical os, potentially improving sperm access to the uterine cavity. These insights bridge biomechanics with reproductive health, offering new avenues for fertility counseling and research. However, further multidisciplinary studies are required to establish evidence-based clinical recommendations.

Acknowledgment

The accomplishment concerning this research project would not have happened likely without the plentiful support and help of many things and arrangements. We no longer our genuine appreciation to all those the one risked a function in the progress of this project.

We would like to express our straightforward recognition to our advisers, Naweed Imam Syed, Professor in the Department of Cell

Biology at the University of Calgary, and Dr. Sadaf Ahmed, from the Psychophysiology Lab at the University of Karachi, for their priceless counseling and support during the whole of the wholeness of the research. Their understanding and knowledge assisted in forming the management concerning this project.

## Declaration of Interest

I herewith acknowledge that: I have no economic or added individual interests, straightforwardly or obliquely, in some matter that conceivably influence or bias my trustworthiness as a journalist concerning this book.

## Conflicts of Interest

The authors profess that they have no conflicts of interest to reveal.

## Financial Support and Protection

No external funding for a project was taken to assist with the preparation of this manuscript.

## References

1. Suarez SS, Pacey AA. Sperm transport in the female generative area. *Hum Reprod Update*. 2006; 12(1): 23–37.
2. Henkel R, Schill WB. Sperm readiness for ART. *Andrologia*. 2003; 35(5): 249–257. (review context).
3. Schultz WW, Camper Andel P, Sabelis I, Mooyaart E. Magnetic reverberation image of male and female organs all the while coitus and female intercourse stimulus. *BMJ*. 1999; 319(7225): 1596–1600. doi:10.1136/bmj.319.7225.1596.
4. Pretorius ES, Timbers K, Malamud D, Barnhart K. Magnetic reverberation is used to decide the distribution of a vaginal coagulate: before, all along, and following in position or time two together simulated and evident communication. *Contraception*. 2002; 66(6): 443–451. doi:10.1016/S0010-7824(02)00412-2.
5. Diemer J, Hahn J, Brandt S, and others. Sperm shift in the organ tract—In silico experiments label key determinants for generative fame. *PLoS Comput Biol*. 2021; 17(7): e1009109. doi:10.1371/chronicle.pcbi.1009109.
6. Nassir M, Levi M, Shaked NT. Dynamic 3D modeling for human semen action through the female cervical waterway and uterine cavity to forecast sperm chance of arriving at the oocyte. *Cells*. 2023; 12(1): 203. Doi:10.3390/cells12010203.
7. Konishi S, Sotome TT, Shimizu K, Oba MS, O'Connor KA, and others. Coital repetitiveness and the chance of gestation in couples trying to understand their first toddler: an anticipated disciple study in Japan. *Int J Environ Res Public Health*. 2020 Jul 10; 17(14): 4985. Doi:10.3390/ijerph17144985. (PMCID: PMC7399901)
8. Mortimer D. The functional study of plants and the study of plants of semen transport. *Hum Reprod*. 1994; 9(12): 2357–2363. (classic tradition).
9. Suarez SS. Regulation of semen storage and transport. *Reproduction*. 2008; 136(5): 615–624.
10. Vehicle Andel P, camper Onna M, de Jong J, Renier WO. Magnetic reverberation depicts male and female organs during sex and female intercourse sexual excitement. *BMJ*. 1999; 319: 1596–1600.
11. Dunphy BC. At sex: beginning deposition and semen introduction. *J Reprod Fertile*. 1987; 81(2): 483–492.
12. Anderson DJ, McNeilly AS. Spermatozoa and cervical mucus interact. *Hum Reprod*. 1984; (classic study).
13. Suarez SS, Pacey AA. The narrative journey of semen through the female generative tract. *Anim Reprod Sci*. 2018.
14. Rudiger M, Cone RA. The popular and mysterious about female generative lot muck. *Bioessays*. 2024; (review on mucus rheology and semen interplay).
15. Mortimer ST. Human spermatozoa: their study of animal and behaviour in cervical gel waste. *Fertil Steril*. 1992; 57(3): 527–535.
16. Suarez SS. Sperm action and the rheological environment: associations for semen pick. *Reprod Fertil Dev*. 2016; 28(1-2): 1–10.
17. Farhi J, Chazan D, Chazan S, and others. Semen dethroning site and gestation rates: cervical against the method of inducing pregnancy. *Fertil Steril*. 2018; (approximate effects).
18. Practice Committee of the American Society for Reproductive Medicine. Optimizing natural pregnancy: a cabinet belief. *Fertil Steril*. 2022; 118(1): 1–12.
19. De Jonge C, Barratt CLR. The search for gametes and fetuses aims to better ART outcomes. *Hum Reprod Update*. 2019; 25(5): 540–555.
20. Van Andel P, Schultz WW, Sabelis I, and others. Sonographic and endoscopic observations of copulation: suggestions for mechanical details of infiltration. *J Sex Med*. 2005; (sonographic coital mechanical details).
21. Weijmar Schultz W, van Andel P, Sabelis I, and others. MRI image of communication: physical changes all along coitus. *J Sex Med*. 2000; (image evidence).
22. Coital biomechanics: displaying pelvic arithmetic and contact forces. *Biomech Mod Mechanobiol*. 2019; (limited item and modeling studies).
23. Andersson K, and others. Coital positions and clitoral anatomy flow: sonographic study. *Sexologies*. 2022; 31(4): 423–430.
24. Post-coital distribution of beginning: ultrasound and MRI studies. *Hum Reprod*. 2001; (disposal image).
25. WebMD. The reality about intercourse positions and getting significant. WebMD. 2023. (services-an architectural finish review rehashing evidence).
26. ASRM practice counseling: timing and commonness of communication vs position in birth advising. ASRM Committee Opinion 2022.
27. Clinical reviews show no strong evidence from two points: coital position and fecundability: diversified practical studies. *Fertil Steril*. 2015–2022.
28. Seminal red body fluid effects on cervical transportation and semen endurance: Rams and ewes models and translational suggestions. *Theriogenology*. 2014; (experimental evidence).
29. Microgravity and inertial belongings on semen action: evidence from fake microgravity experiments. *Reproduction*. 2021; (suggestions for gravity belongings on tiny mobile containers).
30. Methodological pieces of advice: combining image, computational fluid dynamics, and regulated follower designs to judge ejaculate dethroning and semen transport. *J Biomech*. 2020.