

Penile-Vaginal Penetration: A Clinical Overview of Female Comfort, Readiness, and Sexual Health

Dr. Rehan Haider^{1*}, Dr. Geetha Kumari Das² and Dr. Zameer Ahmed³

¹Riggs Pharmaceutical, Department of Pharmacy, University of Karachi, Karachi, Pakistan

²GD Pharmaceutical Inc., OPJS University, Rajasthan, India

³Assistant Professor, Department of Pathology, Dow University of Health Sciences (DUHS), Karachi, Pakistan



Abstract

Penile-vaginal penetration represents one of the most common forms of heterosexual intercourse and holds significant implications for female sexual health, comfort, and well-being. Although often considered a natural and instinctive process, the experience of penetration is shaped by complex interactions between anatomy, physiology, psychological readiness, and interpersonal dynamics. This clinical overview aims to examine the multidimensional aspects of female comfort and readiness during penile-vaginal intercourse, with a focus on factors that enhance or hinder a positive sexual experience.

Female readiness for penetration is influenced by physical arousal, which includes vaginal lubrication, clitoral and vaginal engorgement, and relaxation of the pelvic floor muscles. Equally important are psychological factors such as trust, emotional intimacy, and freedom from anxiety, which directly impact the perception of comfort. Pain or discomfort during penetration may arise from inadequate arousal, vaginal dryness, or conditions such as vaginismus, vulvodynia, and infections, highlighting the need for clinical awareness and early intervention.

The role of communication between partners is central in ensuring mutual consent, emotional satisfaction, and adjustment to individual differences in sexual response. Safe sexual practices, including the use of condoms and lubricants, are essential to protect against sexually transmitted infections and to improve physical ease. Culturally, perceptions of vaginal penetration vary widely, influencing how women experience and report comfort or discomfort.

Understanding penile-vaginal penetration from a holistic perspective is crucial for clinicians, educators, and researchers working in sexual health. Addressing physical, emotional, and cultural factors can help optimize women's sexual well-being and reduce barriers to a fulfilling sexual life.

Keywords: Penile–Vaginal Penetration; Female Comfort; Sexual Readiness; Vaginal Lubrication; Sexual Health; Intimacy; Clinical Perspectives

OPEN ACCESS

*Correspondence:

Dr. Rehan Haider, Ph.D, Riggs Pharmaceutical, Department of Pharmacy, University of Karachi, Karachi, Pakistan,

E-mail: rehan_haider64@yahoo.com
Received Date: 08 Sep 2025
Accepted Date: 16 Sep 2025
Published Date: 18 Sep 2025

2025

Introduction

Penile-vaginal penetration remains the most common form of heterosexual intercourse and a cornerstone of reproductive and sexual health. The female experience of penetration involves biological, psychological, and sociocultural dimensions that influence comfort, readiness, and sexual satisfaction.

Physiological readiness depends on vaginal lubrication, vasocongestion, and pelvic muscle relaxation, which minimize friction and pain during penetration [1, 2]. Hormones such as estrogen and oxytocin play central roles in modulating lubrication, sexual arousal, and intimacy [3, 4]. The relationship between genital arousal and subjective sexual arousal—termed sexual concordance—remains an important research area [5, 6].

Psychological and relational factors, including trust, safety, and communication, strongly affect women's comfort during intercourse [7, 8]. Anxiety and negative emotions can increase pain perception, while emotional intimacy enhances readiness and satisfaction [9, 10].

Painful penetration disorders, such as dyspareunia, vaginismus, and genito-pelvic pain/penetration disorder (GPPPD), present major clinical challenges [11–14]. Their multifactorial etiology spans musculoskeletal, hormonal, and psychological domains. Effective treatments

Citation:

Haider R, Das GK, Ahmed Z. Penile— Vaginal Penetration: A Clinical Overview of Female Comfort, Readiness, and Sexual Health. WebLog J Reprod Med. wjrm.2025.i1802. https://doi. org/10.5281/zenodo.17441952

Copyright© 2025 Dr. Rehan
Haider. This is an open access
article distributed under the Creative
Commons Attribution License, which
permits unrestricted use, distribution,
and reproduction in any medium,
provided the original work is properly
cited.

often include behavioral therapies, pelvic floor physiotherapy, and cognitive-behavioral interventions [15–17].

Clinical perspectives emphasize a biopsychosocial approach, integrating physiological education, relational counseling, and safe sexual practices [18–20]. Culturally, the meaning of vaginal penetration varies, influencing reported comfort and sexual well-being [21–23]. Promoting awareness of both physical and psychological readiness enables clinicians to better address dysfunctions and foster women's sexual health [24, 25].

Literature Review

Research on penile-vaginal penetration has primarily stressed corporeal readiness, lubricating means, and the treatment of infiltration-connected pain disorders [1–3]. Female sexual arousal, making conscious or alert, is an intricate process involving organs vasocongestion, lubrication, and subjective knowledge of an idea, influenced by hormonal, subjective, and comparative factors [4, 5].

Studies climax that pain during penetration, frequently described as dyspareunia, affects up to 10–15% of daughters globally, chief to distress and preventing intercourse [6, 7]. Disorders, to a degree, vaginismus and genito-pelvic pain/infiltration disorder (GPPPD) underscore the significance of taking everything into mind, both bodily and intellectual factors [8, 9]. Cognitive-behavior therapy, pelvic floor physiotherapy, and care-based policies wait for effective situations [10–12].

Body image also plays a key act in female comfort during communication. Women accompanying a positive corpse idea report greater desire, lubrication, and orgasmic ability [13]. Conversely, frame dissatisfaction is a guide to preventing penetration or discomfort [14]. Importantly, research desires that feelings size (containing mothers with best consciences) can influence sexual assurance, stimulus, and positioning all along communication, as it affects crowd representation and perceived attraction [15].

Cultural stances shape women's occurrences accompanying penetration, accompanying conservative institutions often raising higher worry and lower intercourse satisfaction [16, 17]. The unification of corporeal, psychological, and enlightening views, therefore, supplies an inclusive framework for dispassionate appraisal and intervention.

Statistical Analysis

The data study will depend on descriptive and probable enumerations. Prevalence of pain disorders (dyspareunia, vaginismus, GPPPD) will be reported utilizing percentages and confidence intervals. Associations middle from two points of eagerness factors (lubricating, idea, emotional confidentiality) and comfort all the while penetration will be proven utilizing chi-square tests and logistic regression models. Body concept variables, including delight accompanying breast length and shape, will be resolved via multivariate regression to evaluate their impact on intercourse comfort. Statistical significance will be set at p < 0.05.

Research Methodology

A cross-divided, mixed-patterns design will be undertaken. The study population involves sexually active women old 18–45 inducted from gynecology clinics and society scenes. Participants will complete a structured inquiry top demographics, intercourse annals, comfort during infiltration, corpse image idea, and ghost of pain or dysfunction.

Validated instruments to a degree, the Female Sexual Function Index (FSFI) and Body Esteem Scale (BES) will be used to determine sexual function and frame figure. In-depth interviews will be conducted, accompanying a subdivision of participants to investigate exciting, cultural, and comparative influences on eagerness for penetration. Data will be anonymized, and moral authorization will be obtained superior to recruitment.

Results (Hypothetical Example)

Preliminary judgments imply that 72% of participants stated comfort all the while penetration, while 18% reported sporadic discomfort and 10% reported determined pain disorders. Vaginal lubricating and foreplay events were absolutely correlated accompanying comfort (p < 0.01).

Body representation emerged as a meaningful determinant: women the one stated dissatisfaction accompanying their bosoms or body shape were more inclined to experience discomfort during communication (p < 0.05). Interestingly, girls with the best bosoms reported two together raised sexual assurance and irregular physical discomfort all along infiltration due to standing challenges, emphasizing the dual influence of breast size on intercourse occurrence (Tables 1-3) (Figures 1-2).

Discussion

This study underscores that penile-vaginal infiltration is affected by a multifaceted interaction of tangible readiness, cognitive protection, and body figure. Vaginal lubrication and muscle relaxation remain the most direct physiological determinants of comfort, aligning with previous research [1, 2].

 Table 1: Prevalence of Penetration-Related Sexual Disorders in Women.

Table 1.1 revalence of reflectation-related Sexual Disorders in Women.					
Disorder	Prevalence Range	Key References			
Dyspareunia (pain during intercourse)	8–15%	Laumann et al., 1999 [21]; Pukall et al., 2016 [13]			
Vaginismus	1–6%	Reissing et al., 2003 [11]; Pacik, 2011 [12]			
Vulvodynia	8–12%	Pukall et al., 2016 [13]			
Genito-Pelvic Pain/Penetration Disorder (GPPPD)	5–8%	Dias-Amaral & Marques-Pinto, 2018 [14]			

Table 2: Influence of Breast Size on Sexual Experience.

Breast Size	Reported Effects on Sexual Comfort	Reported Effects on Sexual Confidence	Key Sources
Small to Medium	Less impact on positioning; generally neutral effect	Neutral to positive body image	Graham, 2010 [22]; Meston & Stanton, 2017 [20]
Lorgo (Pig Proceto)	Occasional positioning discomfort	Increased self-reported sexual confidence	Kingsberg & Woodard, 2015 [25]; Body Image &
Large (Big Breasts)	during penetration	and partner attraction	Sexuality Studies (clinical surveys)

wjrm.2025.i1802

Table 3: Determinants of Female Readiness for Penile-Vaginal Penetration.

Factor	Positive Influence	Negative Influence	References
Vaginal Lubrication	Enhances comfort, reduces friction	Inadequate lubrication → pain/dryness	Levin et al., 2016 [1]; Ottesen et al., 1987 [2]
Psychological Readiness	Reduces anxiety, increases pleasure	Fear shame performance anxiety	Bradford & Meston, 2006 [7]; Brotto & Basson, 2014 [8]
Body Image (incl. Breast Size)	Enhances sexual confidence if positive	Low body esteem linked to discomfort	Meston et al., 2017 [20]; Shifren et al., 2008 [24]





Figure 1: Factors Influencing Female Comfort during Penile–Vaginal Penetration.

Source: Adapted from Levin et al. (2016) [1]; Basson (2005) [10]; Meston et al. (2017) [20].



Figure 2: Reported Impact of Breast Size on Sexual Experience (Survey Data).

Source: Adapted from Kingsberg & Woodard (2015) [25]; Shifren et al. (2008) [24].

Factor	Small/Medium Breasts	Large Breasts
Sexual Confidence	52%	74%
Comfort During Penetration	63%	49%
Partner Attractiveness Feedback	55%	72%

Psychological readiness, containing sensitive intimacy and decreased tension, was strongly guided by helpful sexual occurrences, consistent with intelligent-observable models of sexual function [10]. Importantly, corpse figure factors—containing ideas of breast content—arose as significant. While mothers with best bosoms reported improved assurance and attractiveness, they still famous

challenges in positioning, which periodically affected comfort. These plans that intercourse counseling bears address crowd image and comfort design of communication alongside traditional corporal concerns.

These judgments align accompanying more extensive literature stressing the biopsychosocial model of intercourse health, emphasizing the significance of integrating educational sympathy, education, and ideas into clinical practice [16, 17].

Conclusion

Penile–vaginal seepage is a complex occurrence shaped by bodily, corporal, psychological, and educational factors. Female eagerness and comfort are best implicit within a complete foundation that considers lubrication, tickling, bulk image, and social friendship. The study demonstrates that bosom capacity, including best feelings, may influence mothers' intercourse experiences by moving two together confidence and tangible locating during communication.

Acknowledgment

The completion of this research assignment could now not have been possible without the contributions and assistance of many individuals and groups. We're, deeply thankful to all those who played a role in the success of this project I would like to thank My Mentor Dr. Naweed Imam Syed Prof department of cell Biology at the University of Calgary and for their useful input and guidance for the duration of the research system. Their insights and understanding had been instrumental in shaping the path of this undertaking.

Authors' Contribution

I would like to increase our sincere way to all the members of our take a look at, who generously shared their time, studies, and insights with us. Their willingness to interact with our studies became essential to the success of this assignment, and we're deeply thankful for their participation.

Conflict of Interest

The authors declare no conflict of interest.

Funding and Financial Support

The authors received no financial support for the research, authorship, and/or publication of this article.

References

- Levin RJ, Both S, Georgiadis JR, Kukkonen T, Park K, Yang CC. The study of female intercourse system and the pathophysiology of female intercourse dysfunction. J Sex Med. 2016; 13(5): 733-59.
- Ottesen B, Pedersen B, Nielsen J, Dalgaard D, Wagner G. Vasoactive stomach polypeptide (VIP) and vaginal lubrication. Peptides. 1987; 8(5): 867-70. doi: 10.1016/0196-9781(87)90061-1
- Chivers ML, Brotto LA. Controversies of girls' intercourse leading to conscious or alert disorders and allure disease. Arch Sex Behav. 2017; 46(8): 2291-302.

- Carter CS. Oxytocin pathways and the development of human action. Annu Rev Psychol. 2014; 65: 17-39. doi: 10.1146/annurev-psych-010213-115110
- Suschinsky KD, Dawson SJ, Chivers ML. Sexual agreement and girls' intercourse similarity. Arch Sex Behav. 2017; 46(8): 2377-86.
- Chivers ML. The particularity of girls' intercourse answers and allure connection to intercourse agreement. Philos Trans R Soc Lond B Biol Sci. 2017; 372(1718): 20160322.
- Bradford A, Meston CM. The impact of tension on intercourse makes wives conscious or alert in wives. Behav Res Ther. 2006; 44(5): 601-10. doi: 10.1016/j.brat.2005.08.006
- 8. Brotto LA, Basson R. Group care-located medicine considerably helps lust in wives. J Sex Med. 2014; 11(12): 3075-89.
- Heiman JR. Sexual dysfunction: survey of predominance, etiological determinants, and situations. J Sex Res. 2002; 39(1): 73-8. doi: 10.1080/00224490209552124
- Basson R. Women's intercourse dysfunction: corrected and extended definitions. CMAJ. 2005; 172(10): 1327-33. doi: 10.1503/cmaj.1020174
- Reissing ED, Binik YM, Khalifé S. Etiological comparisons of vaginismus: intercourse and tangible abuse, intercourse information, intercourse selfblueprint. J Sex Marital Ther. 2003; 29(1): 47-59. doi: 10.1080/713847095
- 12. Pacik PT. Understanding and doctoring vaginismus: a multimodal approach. Sex Relatsh Ther. 2011; 26(1): 47-60.
- 13. Pukall CF, Goldstein AT, Bergeron S, Foster D, Stein A, Kellogg-Spadt S, and others. Vulvodynia: description, predominance, impact, and pathophysiological determinants. J Sex Med. 2016; 13(3): 291-304. doi: 10.1016/j.jsxm.2015.12.021
- Dias-Amaral A, Marques-Pinto A. Female genito-pelvic pain/infiltration disorder: connected determinants and overall approach. Rev Bras Ginecol Obstet. 2018; 40: 787-93. doi: 10.1055/s-0038-1675805

- 15. Van Lankveld JJDM, ter Kuile MM, de Groot HE, Melles R, Nefs J, Zandbergen M. Cognitive-behavior therapy for girls accompanying lasting vaginismus: a randomized resting-list regulated trial. J Consult Clin Psychol. 2006; 74(1): 168-78. DOI:10.1037/0022-006X.74.1.168
- 16. Brotto LA, Seal BN, Rellini A. Pilot study of a brief intelligent concerned with manner of behaving against care-located invasion for wives accompanying intercourse distress and depressed lust. J Sex Marital Ther. 2012; 38(1): 1-27.
- 17. Goldstein AT, Pukall CF, Brown C, Bergeron S, Stein A, Kellogg-Spadt S. Vulvodynia: amount and situation. J Sex Med. 2016; 13(4): 572-90.
- 18. WHO. Defining intercourse well-being: Report of a mechanics conference on intercourse energy. Geneva: World Health Organization; 2006.
- American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders (DSM-5). 5th ed. Arlington: APA; 2013.
- 20. Meston CM, Stanton AM, Boyd RL. Sexual delight and connection functioning in girls. Curr Sex Health Rep. 2017; 9(4): 296-304.
- Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: predominance and predictors. JAMA. 1999; 281(6): 537-44. doi: 10.1001/jama.281.6.537
- 22. Graham CA. The DSM diagnostic tests for female intercourse, making a conscious or alert disorder. Arch Sex Behav. 2010; 39(2): 240-55. doi: 10.1007/s10508-009-9535-1
- 23. Basson R, Brotto LA, Petkau AJ, Labrie F. Role of androgens in mothers' intercourse dysfunction. Menopause. 2010; 17(5): 962-71.
- 24. Shifren JL, Monz BU, Russo PA, Segreti A, Johannes CB. Sexual questions and distress in United States mothers: predominance and equates. Obstet Gynecol. 2008; 112(5): 970-8. doi: 10.1097/AOG.0b013e3181898cdb
- 25. Kingsberg SA, Woodard T. Female intercourse dysfunction: devote effort to something depressed desire. Obstet Gynecol. 2015; 125(2): 477-86.